

HYPERBOLIC MANIFOLDS WHOSE ENVELOPES OF HOLOMORPHY ARE NOT HYPERBOLIC

L. GEATTI, A. IANNUZZI, AND J.-J. LOEB

ABSTRACT. We present a family of examples of two dimensional, hyperbolic complex manifolds whose envelopes of holomorphy are not hyperbolic.

In this note we present a family of hyperbolic complex manifolds whose envelopes of holomorphy are not hyperbolic. Here we consider hyperbolicity in the sense of Kobayashi. In Isaev's classification [2] of two-dimensional hyperbolic manifolds with three-dimensional automorphism group, these manifolds appear as subdomains of $\Delta \times \mathbb{P}^1$, with Δ the unit disk in $\mathbb{C}$ and $\mathbb{P}^1$ the one-dimensional complex projective space. Here we consider their realizations as $SU(1, 1)$ -invariant domains in the Lie group complexification $SL(2, \mathbb{C})/U(1)^{\mathbb{C}}$ of the symmetric space $SU(1, 1)/U(1)$. Then, by applying the univalence results for Stein, equivariant Riemann domains over Lie group complexifications of rank-one, Riemannian symmetric spaces obtained in [1], we can explicitly determine their envelopes of holomorphy. Such envelopes turn out to be all biholomorphic to $\Delta \times \mathbb{C}$. In particular, they are not hyperbolic.

Let G be the Lie group $SU(1, 1)$. Consider the holomorphic action of its universal complexification $G^{\mathbb{C}} = SL(2, \mathbb{C})$ on $\mathbb{P}^1 \times \mathbb{P}^1$ defined by

$$(1) \quad g \cdot ([z_1 : z_2], [w_1 : w_2]) := (g \cdot [z_1 : z_2], \overline{\sigma(g)} \cdot [w_1 : w_2]),$$


where $\sigma(g) = I_{1,1} {}^t \bar{g}^{-1} I_{1,1}$ denotes the conjugation of $G^{\mathbb{C}}$ relative to G (here $I_{1,1}$ the diagonal matrix representing the standard hermitian form of signature $(1, 1)$). Denote by K a maximal compact subgroup of G and by $K^{\mathbb{C}}$ its complexification. The quotient $G^{\mathbb{C}}/K^{\mathbb{C}}$ can be identified with the unique open $G^{\mathbb{C}}$ -orbit in $\mathbb{P}^1 \times \mathbb{P}^1$ given by

$$G^{\mathbb{C}} \cdot ([1 : 0], [1 : 0]) = \{ ([z_1 : z_2], [w_1 : w_2]) \in \mathbb{P}^1 \times \mathbb{P}^1 : z_1 w_1 - z_2 w_2 \neq 0 \}.$$

Mathematics Subject Classification (2000): 32Q45, 32D10, 32M05.

Key words: hyperbolic manifold, envelope of holomorphy, symmetric space.

A global slice for the G -action on $G^{\mathbb{C}}/K^{\mathbb{C}}$ by left translations is represented by the following diagram (cf. [1], Sect. 4.1)



All elements in the diagram, except for z_1 , z_2 and z_3 , lie on hypersurface G -orbits. The points $z_1 := ([1 : 0], [1 : 0])$ and $z_3 := ([0 : 1], [0 : 1])$ lie on G -orbits diffeomorphic to the symmetric space G/K . The point $z_2 := ([1 : i], [1 : i])$ lies on a G -orbit diffeomorphic to a pseudo-Riemannian symmetric space of the same dimension as G/K . The slices $\ell_1, \dots, \ell_4$ are defined by

$$\begin{aligned}\ell_1(t) &= ([\cos \frac{\pi}{4}(1-t) : i \sin \frac{\pi}{4}(1-t)], [\cos \frac{\pi}{4}(1-t) : i \sin \frac{\pi}{4}(1-t)]), \\ \ell_3(t) &= ([\cos \frac{\pi}{4}(1+t) : i \sin \frac{\pi}{4}(1+t)], [\cos \frac{\pi}{4}(1+t) : i \sin \frac{\pi}{4}(1+t)]),\end{aligned}$$

for $0 < t < 1$, and by

$$\begin{aligned}\ell_2(s) &= ([e^s : ie^{-s}], [e^{-s} : ie^s]), \\ \ell_4(s) &= ([e^{-s} : ie^s], [e^s : ie^{-s}]),\end{aligned}$$

for $s > 0$. Note that z_2 is the limit point of ℓ_1 , ℓ_2 , ℓ_3 and ℓ_4 for values of the parameters approaching zero. Similarly one has $\ell_1(1) = z_1$ and $\ell_3(1) = z_3$. The points

$$\begin{aligned}w_1 &:= ([1 : 0], [1 : i]), \quad w_2 := ([1 : i], [0 : 1]), \\ w_3 &:= ([1 : i], [1 : 0]), \quad w_4 := ([0 : 1], [1 : i]).\end{aligned}$$

represent the four non-closed hypersurface G -orbits containing $G \cdot z_2$ in their closure.

Consider the family of G -invariant domains in $G^{\mathbb{C}}/K^{\mathbb{C}}$ defined by

$$D_\beta = G \cdot (z_1 \cup \ell_1((0, 1)) \cup w_1 \cup \ell_2((0, \beta))) \quad \text{for } 0 < \beta < \infty.$$

By the classification of Stein, G -invariant domains in $G^{\mathbb{C}}/K^{\mathbb{C}}$ (cf. Thm. 6.1 in [1]), none of the domains D_β is Stein. Each of them contains a unique Levi-flat orbit given by

$$G \cdot w_1 = \{ ([1 : z], [1 : w]) \in \mathbb{P}^1 \times \mathbb{P}^1 : z \in \Delta, w \in \partial\Delta \} \cong \Delta \times \partial\Delta,$$

and a unique totally-real orbit given by

$$G \cdot z_1 = \{ ([1 : z], [1 : \bar{z}]) \in \mathbb{P}^1 \times \mathbb{P}^1 : z \in \Delta \}.$$

Denote by W the limit domain

$$W = G \cdot (z_1 \cup \ell_1(0, 1) \cup w_1 \cup \ell_2(0, \infty)).$$

By construction W contains every domain D_β . One can check that W coincides with the domain

$$\{ ([1 : z], [w_1 : w_2]) \in \mathbb{P}^1 \times \mathbb{P}^1 : z \in \Delta, w_1 - zw_2 \neq 0 \}.$$

It follows that W is biholomorphic to $\Delta \times \mathbb{C}$ via the map

$$\Delta \times \mathbb{C} \rightarrow W, \quad (u, v) \rightarrow ([1 : u], [1 + uv : v]).$$

In particular, it is Stein and is not hyperbolic.

Proposition 0.1. *For every $0 < \beta < \infty$ the G -invariant domain D_β is hyperbolic. The envelope of holomorphy of D_β is the domain W , which is not hyperbolic.*

Proof. The fact that D_β is hyperbolic follows from Isaev's classification of two-dimensional hyperbolic manifolds with three dimensional automorphism group. Indeed note that for $g \in G$ one has $\overline{\sigma(g)} = \bar{g}$. Thus the restriction to G of the $G^\mathbb{C}$ -action defined in (1) agrees with the G -action given in [2], p.22. As a consequence, D_β coincides with an element of the family denoted there by

$$\widehat{\mathfrak{D}}_t^{(1)} \quad \text{for } 1 < t < \infty$$

and it is hyperbolic.

Denote by $E(D_\beta)$ the envelope of holomorphy of D_β . Since $G^\mathbb{C}/K^\mathbb{C}$ is Stein, the inclusion of D_β in $G^\mathbb{C}/K^\mathbb{C}$ extends to a local biholomorphism p from $E(D_\beta)$ to $G^\mathbb{C}/K^\mathbb{C}$ (cf. [3]). Note that the center $\Gamma = \{\pm Id_2\}$ of G acts ineffectively on $G^\mathbb{C}/K^\mathbb{C}$ and that G/Γ is isomorphic to $SO_0(2, 1)$. Then the $SO_0(2, 1)$ -action on D_β extends to $E(D_\beta)$ and the map p is equivariant, i.e. $p: E(D_\beta) \rightarrow G^\mathbb{C}/K^\mathbb{C}$ is a Stein, $SO_0(2, 1)$ -equivariant Riemann domain.

By Theorem 7.5 in [1], the map p is necessarily injective. Hence $E(D_\beta)$ coincides with the smallest, Stein, G -invariant domain in $G^\mathbb{C}/K^\mathbb{C}$ containing D_β . Since each D_β contains the Levi-flat orbit $G \cdot w_1$, from the classification of Stein, G -invariant domains in $G^\mathbb{C}/K^\mathbb{C}$ given in Theorem 6.1 of [1], it follows that the envelope of holomorphy of D_β is the domain W .

□

Remark. Similarly one can show that W is the envelope of holomorphy of every element in the family of hyperbolic, G -invariant subdomains of W denoted in [2], p. 22, by

$$\mathfrak{D}_{s,t}^{(1)} \quad \text{for} \quad -1 \leq s < 1 < t \leq \infty,$$

where $s = -1$ and $t = \infty$ do not hold simultaneously. In terms of diagram (2) the elements of the family $\mathfrak{D}_{s,t}^{(1)}$ correspond to the domains

$$G \cdot (\ell_1((0, \alpha)) \cup w_1 \cup \ell_2((0, \beta))) \quad \text{for} \quad 0 < \alpha \leq 1 \quad \text{and} \quad 0 < \beta \leq \infty.$$

REFERENCES

- [1] L. GEATTI, A. IANNUZZI, *On univalence of equivariant Riemann domains over a complex semisimple Lie group*. arXiv: math.CV/0612169
- [2] A. V. ISAEV, *Hyperbolic 2-Dimensional Manifolds with 3-Dimensional Automorphism Group*. arXiv: math.CV/0509030 v6
- [3] H. ROSSI, *On envelopes of holomorphy*. Commun. Pure Appl. Math. **16** (1963) 9–17

LAURA GEATTI AND ANDREA IANNUZZI: DIP. DI MATEMATICA, II UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, I-00133 ROMA, ITALY.

E-mail address: geatti@mat.uniroma2.it, iannuzzi@mat.uniroma2.it

JEAN-JACQUES LOEB, DÉPARTEMENT DE MATHÉMATIQUES ET INFORMATIQUES, UNIVERSITÉ D’ANGERS, 2 BOULEVARD LAVOISIER, 49045 ANGERS CEDEX 01

E-mail address: Jean-Jacques.Loeb@univ-angers.fr